



Cite this: DOI: 10.1039/c4nj01904k

Studies of the complexation behavior of tetramorpholinyllo-PNP-lariat ether with Ag(I), Ca(II), Cd(II), Cu(II) and Pb(II) using Electrospray Ionization Mass Spectrometry

Natalia Gutowska,^{*a} Beata Pasternak,^b Piotr Seliger^a and Grzegorz Andrijewski^a

In this publication the cationic metal complexes of tetramorpholinyllo-PNP-lariat ether have been studied using electrospray ionization mass spectrometry (ESI-MS). The tandem mass spectra (MS/MS) of these complexes have also been tested to evaluate the stability of the different types of the complexes formed. As occurred, all selected metal cations form the complexes with 1:1 stoichiometry with the investigated ligand. Only silver ions create a “sandwich” type complex. Furthermore, the divalent cations form complexes with the nitrate anion adduct. In the case of Ca(II), Cd(II) and Pb(II) we also observed another type of the species with an additional water molecule attached to the parent complex.

Received (in Victoria, Australia)
27th October 2014,
Accepted 16th December 2014

DOI: 10.1039/c4nj01904k

www.rsc.org/njc

Introduction

Electrospray ionization mass spectrometry (ESI-MS)^{1–4} is currently an effective tool for analysis of a wide variety of noncovalent complexes,^{5–16} such as those formed in host–guest chemistry. Many studies show that equilibrium distribution of complexes in solution is reflected in the intensities of host–guest complexes observed in the ESI mass spectra.^{17–23} ESI-MS analysis enables us to directly determine the complexes formed in the solution, as well as to study the equilibrium states, carry out quantitative analysis of stability constants or evaluate the binding selectivities.^{24–39} This method is considered to be a “soft ionization” process (involving the transfer of solution ions into the gas phase), and therefore typically yields molecular ions with little or no fragmentation at all. The simplicity of spectra obtained using this technique is of great benefit. Conventional techniques such as infrared spectroscopy, X-ray diffraction and NMR have several drawbacks. The use of ESI-MS for studying metal–ligand interactions and metal complexes is readily recognized. MS is more sensitive than conventional techniques and opens up the possibility of screening the complex samples. Moreover, MS enables simultaneous monitoring of the response of ligand-exchange reactions, additionally obtaining chemical information about the specific compound. Furthermore, ions generated using the electrospray ionization process can be easily fragmented using techniques such as tandem mass

spectrometry (MS/MS). In this method, an ion (called the precursor ion) is selected on the basis of the first stage of MS measurements and activated to produce fragment ions which are then analysed in the second stage of MS.^{40,41} ESI-MS can provide important information concerning the structure, stoichiometry, and metal oxidation state of dissolved metal complexes.^{42,43}

Macrocyclic ligands built on a cyclotriphosphazene ring are compounds which bind cyclophosphazene chemistry with macrocyclic chemistry. The combination of these two fields gives the compounds which have a high possibility for the modification of their structure through the nucleophilic substitution reaction, polycondensation or creation of possible binding centres for cations, anions or neutral species. In this way, the design of various compounds with potential practical use can be realized. Lariat ethers are derived from the group of crown ethers having a side chain attached to the crown moiety by the so-called pivot atoms (C, N or P). The side arms contain atoms or groups with a lone pair of electrons that can cooperate with the heteroatom's electrons from the macrocyclic ring, thus providing a three-dimensional coordination of the guest cation.^{44–46}

Three types of lariat ethers are known depending on the atom to which the side arm is attached: C- (carbon-pivot lariat ethers),^{47–49} N- (nitrogen-pivot lariat ethers)^{50,51} or P- (phosphorus-pivot lariat ethers).⁵² The object of the presented publications is the P-pivot lariat ether derivative. This compound is derived from the group of reactive crown ethers which was formed by incorporation of a chloro-substituted cyclotriphosphazene unit into the macrocyclic polyether skeleton to give mono- and bis-cyclosubstituted derivatives.⁵³ By substitution of the reactive chloride substituents

^a Department of Inorganic and Analytical Chemistry, Faculty of Chemistry, University of Łódź, Tamka 12, 91-403 Łódź, Poland. E-mail: natalia_gutowska@wp.pl

^b Laboratory of Molecular Spectroscopy, Faculty of Chemistry, University of Łódź, Tamka 12, 91-403 Łódź, Poland






Fig. 1 Investigated ligand L.

with various nucleophiles we can obtain a whole series of PNP-lariat ethers. There are known PNP-lariat ether derivatives substituted with different amines^{54–58} and sodium arylates.^{59,60} These compounds exhibit very good abilities toward complexation. Due to the presence of the binding sites in the form of both oxygen and nitrogen donor atoms, these ligands form stable complexes with alkali metals and alkaline earth metal cations as well as transition metal cations.^{61,62}

Macrocyclic derivatives of cyclotriphosphazenes and their complexes are recognized as compounds of proven antitumor activity⁶³ as well as anti-AIDS activity.⁶⁴ Anti-proliferative activity may be modified by the formation of complexes with metal ions. These types of compounds form selective bindings with silver ions.^{65,66} Additionally, recent investigations showed that silver compounds exhibit antibacterial properties.⁶⁷ It has been proved that the thio-substituted PNP-lariat ethers are commonly used to remove highly toxic heavy metal pollutants and for recovery of silver from industrial wastes.⁶⁸ Complexing abilities of these compounds give the possibility of modelling biological systems, in which sulphur-organic compounds are involved.^{69,70} The PNP-crown amino ethers can be potentially useful as “pH-controlled active ion carriers” in liquid membranes.⁷¹ However, bis-PNP-lariat ethers were found to behave as efficient ion carriers for heavy metal (Zn^{2+} , Cd^{2+} , Pb^{2+}) transport carriers across polymer inclusion membranes, in particular for lead(II), due to the formation of “sandwich” type complexes with the macrocyclic compounds.⁷²

In the present paper we report the ESI-MS studies of the tetramorpholinyl-PNP-lariat ether complexes with $\text{Ca}(\text{II})$, $\text{Ag}(\text{I})$, $\text{Cd}(\text{II})$, $\text{Cu}(\text{II})$ and $\text{Pb}(\text{II})$ ions. The tandem mass spectra (MS/MS)





Fig. 2 The mass spectrum of investigated ligand L.



Scheme 1 Fragmentation pattern of the free ligand.



Table 1 Positive ion ESI-MS data for the complexes

Metal salt	Ions (m/z)
AgNO ₃	[Ag(I)L] ⁺ (778.5) [Ag(I)L ₂] ⁺ (1449.7)
Ca(NO ₃) ₂	[Ca(II)L] ²⁺ (355.7) [Ca(II)L + NO ₃] ⁺ (773.5) [Ca(II)L + OH] ⁺ (728.5)
Cu(NO ₃) ₂	[Cu(I)L] ⁺ (734.6) [Cu(II)L] ²⁺ (367.3) [Cu(II)L + NO ₃] ⁺ (796.5)
Cd(NO ₃) ₂	[Cd(II)L] ²⁺ (392.8) [Cd(II)L + NO ₃] ⁺ (847.6) [Cd(II)L + H ₂ O] ⁺ (802.5)
Pb(NO ₃) ₂	[Pb(II)L] ²⁺ (439.8) [Pb(II)L + NO ₃] ⁺ (941.6) [Pb(II)L + H ₂ O] ⁺ (896.6)

of these complexes were also examined to evaluate the stability of the different complex types. The tetramorpholinyl-PNP-lariat ether has a few structural units with the potential ability to take part in the binding of metal cations: the polyether oxygen donors of the macrocyclic PNP-crown skeleton, the endocyclic nitrogen atom of the cyclotriphosphazene ring and the exocyclic nitrogen atoms and oxygen donor atoms of the morpholinyl substituents. This ligand is capable of complexing “hard” and “soft” cations.

Experimental

Synthesis

The synthesis of tetramorpholinyl-PNP-lariat ether has been previously reported.⁶⁵ The investigated ligand is presented in Fig. 1.

Materials

AgNO₃ (Sigma Aldrich) was used as received. Nitrates of Ca(II), Cd(II), Cu(II) and Pb(II) were purchased from POCh Gliwice and used without further purification.

Methanol (HPLC-grade) was purchased from J.T. Baker (POCh Gliwice) and used without purification.

Stock solutions of the ligand and metals salts in methanol (10⁻³ M each) were prepared prior to dilution to 10⁻⁴ M for mass spectrometric investigation (ligand/metal salt ratio 1 : 1). In each case freshly prepared solutions were used.

Instrumental methods

ESI-MS and MS/MS were recorded using a Varian 500-MS LC ion-trap mass spectrometer (Palo Alto, CA, USA). The whole sample was introduced into ESI-MS source by continuous infusion by means of the instrument syringe pump at a rate of 10 µl min⁻¹. The ESI-source was operated at 5.00 kV and the capillary heater was set to 300 °C. The cone voltage was set within the range of 40–260 V. Scanning was performed from m/z = 100 to 1500. For fragmentation experiments, mass-selected monoisotopic molecular ions were isolated in the ion trap and collisionally activated using helium damping gas present in the mass analyser as a collision gas. The experiments were performed in positive ion-mode.

Results and discussion

The ESI-MS spectrum of L revealed the presence of the protonated molecule, at m/z 672.6, and a less abundant ion corresponding to the Na⁺ adduct of the parent compound Fig. 2. ESI-MS fragmentation analysis of L is shown on Scheme 1. In the first step we observed the loss of morpholinyl substituent (m/z = 585.4), probably from the exomacrocyclic P atom. Such a behaviour for a tetrapyrrolidiny cyclophosphazene derivative was described previously.⁶² There was also whole substituent loss observed without the cleavage of the ring. In the following steps the loss of successive fragments of the polyether chain was noticed. After detachment of the two -OCH₂CH₂ fragments we observed two concurrent fragmentation patterns. The first involved a stepwise loss of one morpholinyl substituent (m/z = 412.2) and the subsequent -OCH₂CH₂ group (m/z = 368.2). The second one showed two major fragment ions with m/z = 453.3 due to the loss of the




Fig. 3 The mass spectrum of L with AgNO₃ (the attached fragment of AgL₂ spectrum had RI < 5%).



–OCH₂CH₂ moiety and with $m/z = 368.2$ caused by the loss of the morpholinyl substituent. In the last stage splitting-off of the third morpholinyl substituent is observed.

The ESI mass spectra of the mixture of L with metal ions (Ca, Ag, Cd, Cu and Pb) indicate the presence of many different forms of coordination compounds. The experiments were done using nitrate salts because the stability constants of the

investigated ligand with silver(i) ions (with nitrate as counter ions) had been reported earlier in the literature.⁶⁵ The resulting complexes were presented in Table 1. As occurs from the analysis of mass spectra – all the metal ions (apart from the calcium ion) are bound with the endocyclic nitrogen atom. The nitrogen atom built-in the structure of the polyether represents a “soft” electron donor place to which the “soft” transition




Fig. 4 Comparison of the theoretical isotope pattern calculation (a) for $[Ag(I)L]^+$ with the one observed experimentally (b).




Fig. 5 The ESI-MS/MS spectrum of ion $[^{107}Ag(I)L]^+$ – $m/z = 778.5$.



metals cations as well as heavy metals cations are preferably bound. The oxygen atoms from crown ether moiety are “hard” donors having the affinity toward “hard” cations of I and II group of the periodic table of elements.⁷³

The mass spectrum of L with AgNO₃ was dominated by the [Ag(I)L]⁺ complex. Among other investigated metal ions only the silver ion form a “sandwich” type complex with two molecules of the ligand (peak at *m/z* = 1449.7). The interesting fact about the system with the Ag(I) ion was no signal of the uncomplexed ligand (no peak at *m/z* = 672.6), what probably may confirm the very strong binding of silver ions by the investigated ligand. The complexes of AgL and AgL₂ type obtained during the ESI-MS measurements are in accordance with the results obtained using the potentiometric method.⁶⁵ Divalent cations form complexes with the nitrate anion or with water molecule attached to parent compound, such behaviour was observed for Cd(II), Pb(II) and Ca(II) ions. For the complexes with divalent cations, the loss of one counter ion (NO₃[−]) seems to be the predominated ionization process. The ESI-MS spectra of L with Cd(II), Cu(II) and Pb(II) show the presence of [Cd(II)L]²⁺, [Cu(II)L]²⁺ and [Pb(II)L]²⁺ complexes. The relative intensity of these complexes was

very low (RI < 5%). The situation is different in the case of calcium ions – because in this case the formed [Ca(II)L]²⁺ complex exhibits much higher relative abundance of 31%.

In the case of the copper ion due to the measurement conditions we observed the reduction process Cu(II) → Cu(I). This type of process is a phenomenon observed in ESI^{74–77} and is usually not an electrochemical redox reaction. This process can be explained as an electron transfer between a Cu complex and a solvent molecule in the gas phase.

In the case of silver ions we observed two types of complexes [Ag(I)L]⁺ and [Ag(I)L₂]⁺ – Fig. 3. The ESI-MS for [Ag(I)L]⁺ show two main peaks at *m/z* 778.5 and 780.5, which are expanded in Fig. 4(b), and compared well with the corresponding calculated (Fig. 4(a)) isotope pattern. The isotope patterns in Fig. 4 are consistent with a molecular cation containing one silver atom and exhibits the natural intensity ratio for silver isotopes.

We made the fragmentation of the peak *m/z* = 778.5. The fragmentation pattern for [Ag(I)L]⁺ shows the same fragmentation pattern as in the case of the ligand Fig. 5. This suggests that binding of silver ions by the ligand molecule is reasonably




Fig. 6 The mass spectrum of L with Ca(NO₃)₂.





Fig. 7 The ESI-MS/MS spectrum of ion [Ca(II)L + NO₃]⁺ – *m/z* = 773.5.



strong and complex formed does not completely disintegrate under the influence of the ionization process.

The ESI-MS of an equimolar mixture of L and $\text{Ca}(\text{NO}_3)_2$ shows three types of complexes: $[\text{Ca}(\text{II})\text{L}]^{2+}$, $[\text{Ca}(\text{II})\text{L} + \text{NO}_3]^+$ and $[\text{Ca}(\text{II})\text{L} + \text{OH}]^+$ (Fig. 6). The most abundant peak at m/z 773.5 is the one corresponding to $[\text{Ca}(\text{II})\text{L} + \text{NO}_3]^+$ complex. The ESI-MS/MS spectrum of $[\text{Ca}(\text{II})\text{L} + \text{NO}_3]^+$ complex is shown in Fig. 7. For this complex we observed the loss of nitrate anion and one proton (m/z = 710.5). The MS/MS fragmentation analysis of calcium complex revealed at the first stage the loss of one morpholinyl substituent (probably from the exomacrocyclic

P atom) and $-(\text{CHCH}_2)$ group (m/z = 597.4). Then we saw an intense peak at m/z 552.5, corresponding to the loss of fragment of the polyether chain $-(\text{OCH}_2\text{CH}_2)$. From this step two concurrent fragmentation paths were observed. In the first one – two less abundant peaks at m/z = 465.4 and m/z = 406.2, corresponding to the loss of second morpholinyl group from exomacrocyclic P atom and another fragment of the crown ether ring and H_2O molecule. The second fragmentation pattern involves a stepwise loss of $-(\text{OCH}_2\text{CH}_2)$ and OH moiety to an ion with m/z 491.4, followed by losses of morpholinyl substituent to an ion with m/z 406.2. These two fragmentation patterns are presented in Scheme 2.



Scheme 2 Fragmentation pattern of calcium(II) complex.




Fig. 8 Comparison of the theoretical isotope pattern calculation (a) for calcium complex at m/z = 406.2 with the experimentally observed pattern (b).







Fig. 9 The ESI-MS/MS spectrum of complex ion $[\text{Ca(II)L} + \text{OH}]^+ - m/z = 728.5$.



Scheme 3 Fragmentation pattern of $[\text{Ca(II)L} + \text{OH}]^+$ complex.




Fig. 10 The mass spectrum of L with $\text{Cd}(\text{NO}_3)_2$ (the attached fragment of CdL spectrum had RI < 5%).

The region within the peak at m/z 406.2, which is expanded in Fig. 8(b), compare well with the corresponding calculated (Fig. 8(a)) isotope pattern.

Additionally the fragmentation of the $[\text{Ca(II)L} + \text{OH}]^+$ complex was checked (Fig. 9). In this case we also noticed two fragmentation patterns. At the beginning we observed the loss






Fig. 11 The ESI-MS/MS spectrum of ion $[\text{Cd}(\text{II})\text{L} + \text{NO}_3]^+ - m/z = 847.6$.




Fig. 12 The mass spectrum of L with $\text{Cu}(\text{NO}_3)_2$ (the attached fragment of $\text{Cu}(\text{II})\text{L}$ spectrum had RI < 5%).





Fig. 13 The ESI-MS/MS spectrum of ion $[\text{Cu}(\text{II})\text{L} + \text{NO}_3]^+ - m/z = 796.5$.

of $-\text{OH}$ fragment as H_2O molecule (18 Da) to an ion with $m/z = 710.5$. The most intense peak m/z 684.5 is connected with the loss of $-(\text{C}_2\text{H}_2)$ group which was followed by the loss of 44 Da $-(\text{OCH}_2\text{CH}_2)$. The less intense signal ($m/z = 666.6$) is originated





Scheme 4 Fragmentation pattern of $[\text{Cu}(\text{II})\text{L} + \text{NO}_3]^+$ complex.

Scheme 5 Fragmentation pattern of $[\text{Cu}(\text{I})\text{L}]^+$ complex.

from the loss of $-(\text{OCH}_2\text{CH}_2)$ and that with m/z 640.4 due to subsequent loss of $-(\text{C}_2\text{H}_2)$. The proposal of the fragmentation pattern is presented in Scheme 3.

The ESI-MS spectra of **L** and $\text{Cd}(\text{NO}_3)_2$ show three types of complexes: $[\text{Cd}(\text{II})\text{L}]^{2+}$, $[\text{Cd}(\text{II})\text{L} + \text{NO}_3]^+$ and $[\text{Cd}(\text{II})\text{L} + \text{H}_2\text{O}]^+$ (Fig. 10). We made the fragmentation of the peak $m/z = 847.6$ and we observed the same fragmentation pattern as in the case of the free ligand Fig. 11. This fact proves that the cadmium(II) complex is very unstable structure. A similar situation was observed during the fragmentation of the $[\text{Cd}(\text{II})\text{L} + \text{H}_2\text{O}]^+$ complex.

The copper ions create three types of complexes: $[\text{Cu}(\text{I})\text{L}]^+$, $[\text{Cu}(\text{II})\text{L}]^{2+}$ and $[\text{Cu}(\text{II})\text{L} + \text{NO}_3]^+$ (Fig. 12). The MS/MS fragmentation pattern for the $[\text{Cu}(\text{II})\text{L} + \text{NO}_3]^+$ complex ($m/z = 796.5$) shows the loss of $-\text{NO}_3$ to an ion with $m/z = 734.6$ (Fig. 13). During the next step the loss of the morpholinyl group with $m/z = 647.6$ was noticed. Then we observed the loss of another morpholinyl substituent together with the $-(\text{CH}_2\text{CH}_2)$ group with $m/z = 534.6$.

In the last step the splitting-off of the fragments of the polyether moiety was observed (Scheme 4). Fragmentation of $[\text{Cu}(\text{I})\text{L}]^+$ with m/z 734.6 (Fig. 14) proceeds with the loss of two morpholinyl substituents ($m/z = 562.4$). The most abundant fragment observed at $m/z = 477.4$ was formed due to the loss of another morpholinyl group. Then the losses of 28 Da $-(\text{CH}_2\text{CH}_2)$ peaks with $m/z = 448.3$ and with $m/z = 363.2$ due to subsequent loss of the last morpholinyl group were observed. This fragmentation pattern is shown on Scheme 5. Analysing the ionic radii of Cu(I) and the probable ligand cavity size we can notice that the copper ion fits well into crown moiety. Crown size (170–220 pm) and copper(I) ion (182 pm)⁷⁸ are comparable, while the copper(II) ion (144 pm) is much smaller hence almost whole fragment of the polyether chain decay is observed.

In the case of lead(II) ions we observed three types of complexes: $[\text{Pb}(\text{II})\text{L}]^{2+}$, $[\text{Pb}(\text{II})\text{L} + \text{NO}_3]^+$ and $[\text{Pb}(\text{II})\text{L} + \text{H}_2\text{O}]^+$ (Fig. 15). The fragmentation pattern of $[\text{Pb}(\text{II})\text{L} + \text{NO}_3]^+$ shows (as in the case of copper derivative) the loss of $-\text{NO}_3$ group to an ion with $m/z = 878.7$. Then the loss of one morpholinyl substituent giving the peak at $m/z = 793.6$ was observed. The most intense peak




Fig. 14 The ESI-MS/MS spectrum of ion $[\text{Cu}(\text{I})\text{L}]^+ - m/z = 734.5$.






Fig. 15 The mass spectrum of L with $\text{Pb}(\text{NO}_3)_2$.





Fig. 16 The ESI-MS/MS spectrum of ion $[\text{Pb}(\text{II})\text{L} + \text{NO}_3]^+ - m/z = 878.7$.



Scheme 6 Fragmentation pattern of $[\text{Pb}(\text{II})\text{L} + \text{NO}_3]^+$ complex.




at $m/z = 765.5$ is related to the fragment which remained after $-(\text{CH}_2\text{CH}_2)$ loss. The further decomposition was realized through the stepwise splitting-off almost all parts of the polyether moiety ($m/z = 633.5$). In the final step the morpholinyl group was detached giving the peak at $m/z = 546.3$ (Fig. 16), (Scheme 6).

Conclusion

Electrospray ionization mass spectrometry was used to probe the complexation behaviour of tetramorpholinyl-PNP-lariat ether with a wide range of metals ions. All selected metal cations form the complexes with 1:1 stoichiometry with the investigated ligand. Only silver ions create the “sandwich” type complex $[\text{Ag}(\text{I})\text{L}_2]^+$. Furthermore divalent cations form complexes with the nitrate anion adduct. In the case of $\text{Ca}(\text{II})$, $\text{Cd}(\text{II})$ and $\text{Pb}(\text{II})$ we noticed also another type of complex species where water molecules are involved in the formation of the molecular ion observed during the measurements: $[\text{Ca}(\text{II})\text{L} + \text{OH}]^+$, $[\text{Cd}(\text{II})\text{L} + \text{H}_2\text{O}]^+$ and $[\text{Pb}(\text{II})\text{L} + \text{H}_2\text{O}]^+$. Additionally, as far as copper ions are concerned, what we observed during the ionisation process was the reduction of $\text{Cu}(\text{II}) \rightarrow \text{Cu}(\text{I})$ and formation of the complexes with both copper(I) and copper(II) ions. As we noticed from the fragmentation pattern, the complex with the cadmium ion is unstable under the measurement conditions. Calcium, copper and lead forms of complexes in the MS/MS fragmentation pattern of $[\text{M}(\text{II})\text{L} + \text{NO}_3]^+$ show the loss of the $-\text{NO}_3$ anion. In the case of tandem MS/MS for all investigated complexes – at the beginning we always observed the loss of the morpholinyl substituent and subsequent splitting-off of the polyether chain. Breaking of the chain and loss of the $-(\text{CH}_2\text{CH}_2)$ and $-(\text{OCH}_2\text{CH}_2)$ group fragments were observed in the literature in case of common crown ethers derivatives.⁷⁹ Only in the case of copper(I) we observed the loss of four morpholinyl substituents. This fact suggests that the copper(I) is strongly accommodated inside the polyether ring. The obtained results suggest that ESI-MS technique seems to be a promising tool for characterization of complexes which are existing in the solution, despite the differences in the phase of the measurement conditions.

Notes and references

- M. Yamashita and J. B. Fenn, *J. Phys. Chem.*, 1984, **88**, 4451.
- R. B. Cole, *J. Mass Spectrom.*, 2000, **35**, 763–772.
- J. B. Fenn, M. Mann, C. K. Meng, S. F. Wong and C. M. Whitehouse, *Mass Spectrom. Rev.*, 1990, **9**, 37–70.
- Electrospray Ionization Mass Spectrometry: Fundamentals, Instrumentation and Applications*, ed. R. B. Cole, Wiley-Interscience, New York, NY, 1997, pp. 1–577.
- J. A. Loo, *Int. J. Mass Spectrom.*, 2000, **200**, 175–186.
- C. A. Schalley, *Int. J. Mass Spectrom.*, 2000, **194**, 11–39.
- J. M. Daniel, S. D. Friess, S. Rajagopalan, S. Wendt and R. Zenobi, *Int. J. Mass Spectrom.*, 2002, **216**, 1–27.
- T. D. Veenstra, *Biophys. Chem.*, 1999, **79**, 63–79.
- P. B. Armentrout, C. A. Austin and M. T. Rodgers, *J. Phys. Chem.*, 2014, **118**, 8088–8097.
- J. S. Brodbelt, *Int. J. Mass Spectrom.*, 2000, **200**, 57–69.
- N. Shen, R. Marshall Pope and D. V. Dearden, *Int. J. Mass Spectrom.*, 2000, **195/196**, 639–652.
- R. Frański and B. Gierczyk, *J. Am. Soc. Mass Spectrom.*, 2010, **21**, 545–549.
- R. Frański, *Rapid Commun. Mass Spectrom.*, 2009, **23**, 3488–3491.
- A. Tsuda, H. Moriwaki and T. Oshima, *J. Chem. Soc., Perkin Trans. 2*, 1999, 1235–1240.
- T. Ly and R. R. Julian, *J. Am. Soc. Mass Spectrom.*, 2006, **17**, 1209–1215.
- S. F. Ralph, M. M. Sheil, L. A. Rodney, J. Geue and A. M. Sargeson, *J. Chem. Soc., Dalton Trans.*, 1996, 4417–4424.
- D.-S. Young, H.-Y. Hung and L. K. Liu, *J. Mass Spectrom.*, 1997, **32**, 432–437.
- S. M. Blair, J. S. Brodbelt, G. M. Reddy and A. P. Marchand, *J. Mass Spectrom.*, 1998, **33**, 721–728.
- E. C. Kempen, J. S. Brodbelt, R. A. Bartsch, Y. Jang and J. S. Kim, *Anal. Chem.*, 1999, **10**, 5494–5500.
- E. C. Kempen, J. S. Brodbelt and M. Reyzer, *Struct. Chem.*, 1999, **10**, 213–220.
- M. Reyzer, J. S. Brodbelt, A. P. Marchand, Z. Chen, Z. Huang and I. N. N. Namboothiri, *Int. J. Mass Spectrom.*, 2001, **200**, 57–69.
- S. M. Blair, J. S. Brodbelt, A. P. Marchand, H.-S. Chong and S. Alidhodzic, *J. Am. Soc. Mass Spectrom.*, 2000, **11**, 884–891.
- V. B. Di Marco and G. G. Bombi, *Mass Spectrom. Rev.*, 2006, **25**, 347–379.
- A. Chapeaurouge, L. Bigler, A. Schafer and S. Bienz, *J. Am. Soc. Mass Spectrom.*, 1995, **6**, 207.
- K. Wang and R. B. Cole, *Org. Mass Spectrom.*, 1994, **29**, 419.
- K. Wang and G. W. Gokel, *J. Org. Chem.*, 1996, **61**, 4693–4697.
- S. M. Blair, J. S. Brodbelt, A. P. Marchand, K. A. Kumar and H. S. Chong, *Anal. Chem.*, 2000, **72**, 2433–2445.
- S. M. Blair, E. C. Kempen and J. S. Brodbelt, *J. Am. Soc. Mass Spectrom.*, 1998, **9**, 1049–1059.
- D.-S. Young, H.-Y. Hung and L. K. Liu, *Rapid Commun. Mass Spectrom.*, 1997, **11**, 769–773.
- E. Leize, A. Jaffrezic and A. Van Dorsselaer, *J. Mass Spectrom.*, 1996, **31**, 537–544.
- S. Williams, S. M. Blair, J. S. Brodbelt, X. Huang and R. A. Bartsch, *Int. J. Mass Spectrom.*, 2001, **212**, 389–401.
- C. L. Scherman and J. S. Brodbelt, *Anal. Chem.*, 2003, **75**, 1828–1836.
- J. B. Nicoll and D. V. Dearden, *Int. J. Mass Spectrom.*, 2001, **204**, 171–183.
- T. Oshima, F. Matsuda, K. Fukushima, H. Tamura, G. Matsubayashi and R. Arakawa, *J. Chem. Soc., Perkin Trans. 2*, 1998, 145–148.
- R. A. Bartsch and M. D. Eley, *Tetrahedron*, 1996, **52**, 8979–8988.
- W. Z. Shou and R. F. Browner, *Anal. Chem.*, 1999, **71**, 2265–3373.
- J. G. Krabbe, A. R. de Boer, G. van der Zwan, H. Lingeman, W. M. A. Niessen and H. Irth, *J. Am. Soc. Mass Spectrom.*, 2007, **18**, 707–713.



- 38 A. Ohki, J. Ping Lu, X. Huang and R. A. Bartsch, *Anal. Chem.*, 1994, **66**, 4332–4336.
- 39 S. M. Williams, J. S. Brodbelt and R. A. Bartsch, *J. Am. Soc. Mass Spectrom.*, 2003, **14**, 1215–1228.
- 40 K. Vékey, *J. Chromatogr. A*, 2001, **921**, 227–236.
- 41 O. Núñez, E. Moyano and M. T. Galceran, *Trends Anal. Chem.*, 2005, **24**, 683–703.
- 42 A. R. S. Ross, M. G. Ikonomon, J. A. Thompson and K. I. Orians, *Anal. Chem.*, 1998, **70**, 2225–2235.
- 43 X. Yu, M. Wojciechowski and C. Fenselau, *Anal. Chem.*, 1993, **65**, 1355–1359.
- 44 G. W. Gokel, *Chem. Soc. Rev.*, 1992, **39**, 39–47.
- 45 G. W. Gokel, *Crown Ethers and Cryptand*, University of Miami Coral Gables, Florida, USA, 1991.
- 46 K. Brandt, I. Porwollik and M. Siwy, *Wiadomości chemiczne, Chemia supramolekularna*, 1997, pp. 91–117.
- 47 D. M. Dishong, C. J. Diamond, M. I. Cinoman and G. W. Gokel, *J. Am. Chem. Soc.*, 1983, **105**, 586.
- 48 D. M. Goli, D. M. Dishong, C. J. Diamond and G. W. Gokel, *Tetrahedron Lett.*, 1982, **23**(50), 5243–5246.
- 49 D. B. White, K. A. Arnold and G. W. Gokel, *Tetrahedron Lett.*, 1987, **28**(16), 1749–1752.
- 50 R. A. Schultz, D. B. White, D. M. Dishong, K. A. Arnold and G. W. Gokel, *J. Am. Chem. Soc.*, 1985, **107**, 6659.
- 51 D. M. Dishong, C. J. Diamond, M. I. Cinoman and G. W. Gokel, *J. Am. Chem. Soc.*, 1983, **105**, 586.
- 52 K. Brandt, T. Kupka, J. Drozd, J. C. van de Grampel, A. Meetsma and A. P. Jekel, *Inorg. Chim. Acta*, 1995, **228**(2), 187–192.
- 53 K. Brandt, I. Porwollik, T. Kupka, R. A. Shaw and D. B. Davies, *J. Org. Chem.*, 1995, **60**, 7433–7438.
- 54 I. Porwollik-Czomperlink, K. Brandt, T. A. Clayton, D. B. Davies, R. J. Eaton and R. A. Shaw, *Inorg. Chem.*, 2002, **41**(19), 4944–4951.
- 55 S. Besli, *Inorg. Chem. Commun.*, 2005, **8**, 449–452.
- 56 K. Brandt, I. Porwollik-Czomperlink, M. Siwy, T. Kupka, R. A. Shaw, D. B. Davies and R. A. Bartsch, *J. Inclusion Phenom. Macrocyclic Chem.*, 1999, **35**, 281–289.
- 57 K. Brandt, I. Porwollik, M. Siwy, T. Kupka, R. A. Shaw, D. B. Davies, M. B. Hursthouse and G. D. Sykara, *J. Am. Chem. Soc.*, 1997, **119**, 1143–1144.
- 58 D. B. Davies, T. A. Clayton, R. J. Eaton, R. A. Shaw, A. Egan, M. B. Hursthouse, G. D. Sykara, I. Porwollik-Czomperlik, M. Siwy and K. Brandt, *J. Am. Chem. Soc.*, 2000, **122**, 12447–12457.
- 59 R. A. Bartsch, E. K. Lee, S. Chun, N. Elkarim, K. Brandt, I. Porwollik-Czomperlink, M. Siwy, D. Lach and J. Silverring, *J. Chem. Soc., Perkin Trans. 2*, 2002, 442–448.
- 60 K. Brandt, I. Porwollik, T. Kupka, A. Olejnik, R. A. Shaw and D. B. Davies, *J. Org. Chem.*, 1995, **60**, 7433–7438.
- 61 P. Seliger, G. Andrijewski, M. Siwy and D. Sęk, *Pol. J. Chem.*, 2009, **83**, 581–588.
- 62 K. Brandt, P. Seliger, A. Grzejdzia, T. J. Bartczak, R. Kruszyński, D. Lach and J. Silberring, *Inorg. Chem.*, 2001, **40**, 3704–3710.
- 63 M. Siwy, D. Sęk, B. Kaczmarczyk, I. Jaroszewicz, A. Nasulewicz, M. Pelczyńska, D. Nevozhay and A. Opolski, *J. Med. Chem.*, 2006, **49**, 806–810.
- 64 K. Brandt, R. Kruszyński, T. J. Bartczak and I. Porwollik-Czomperlink, *Inorg. Chim. Acta*, 2001, **322**, 138–144.
- 65 P. Seliger, N. Sołtys, G. Andrijewski and M. Siwy, *New J. Chem.*, 2012, **36**, 2607–2612.
- 66 N. Sołtys, P. Seliger, G. Andrijewski and M. Siwy, *RSC Adv.*, 2013, **3**, 25351–25359.
- 67 S. A. Cotton, *The Chemistry of Precious Metals*, Blackie Academic & Professional, an Imprint of Chapman & Hall, London UK, 1997, p. 325.
- 68 R. Kruszyński, M. Siwy, I. Porwollik-Czomperlink and A. Trzesowska, *Inorg. Chim. Acta*, 2006, **359**, 649–658.
- 69 T. Nabeshima, *Coord. Chem. Rev.*, 1996, **148**, 151.
- 70 V. V. Litvinova and A. V. Anisimov, *Khim. Geterotsikl. Soedin.*, 1999, 1587.
- 71 M. Penso, A. Maia, V. Lupi and G. Tricarico, *Tetrahedron*, 2011, **67**, 2096–2102.
- 72 C. A. Kozłowski and J. Kozłowska, *J. Membr. Sci.*, 2009, **326**, 215–221.
- 73 in *Crown ethers and analogs*, ed. S. Patai and Z. Rappoport, Wiley, Chichester, 1989.
- 74 L. Gianelli, V. Amendola, L. Fabbri, P. Pallavicini and G. G. Mellerio, *Rapid Commun. Mass Spectrom.*, 2001, **15**, 2347–2353.
- 75 H. Lavanant, H. Virelizier and Y. Hoppilliard, *J. Am. Soc. Mass Spectrom.*, 1998, **9**, 1217–1221.
- 76 D. Schroder, T. Weiske and H. Schwarz, *Int. J. Mass Spectrom.*, 2002, **219**, 729.
- 77 A. Tintaru, L. Charles, P. Milko, J. Roithova and D. Schroder, *J. Phys. Org. Chem.*, 2009, **22**, 229.
- 78 *The Elements*, ed. J. Emsley, Clarendon Press, Oxford, 2 edn, 1991.
- 79 H. Gleispach, H. J. Leis, C. Erk and M. Bulut, *J. Inclusion Phenom. Mol. Recognit. Chem.*, 1996, **24**, 163–173.

